

KEYTRUDA:

Approved for use regardless of BRAF status

KEYTRUDA is indicated for the treatment of patients with unresectable or metastatic melanoma.

Dear Customer:

Merck is pleased to announce that KEYTRUDA is now approved for 200-mg fixed-dose administration in patients with unresectable or metastatic melanoma.



Now approved for 200-mg Q3W fixed-dose administration in advanced melanoma

The recommended dose of KEYTRUDA is 200-mg fixed dose administered as an intravenous infusion over 30 minutes Q3W until disease progression or unacceptable toxicity.

Q3W = every 3 weeks.

SELECTED SAFETY INFORMATION

- KEYTRUDA can cause immune-mediated pneumonitis, including fatal cases. Pneumonitis occurred in 94 (3.4%) of 2799 patients receiving KEYTRUDA, including Grade 1 (0.8%), 2 (1.3%), 3 (0.9%), 4 (0.3%), and 5 (0.1%) pneumonitis, and occurred more frequently in patients with a history of prior thoracic radiation (6.9%) compared to those without (2.9%). Monitor patients for signs and symptoms of pneumonitis. Evaluate suspected pneumonitis with radiographic imaging. Administer corticosteroids for Grade 2 or greater pneumonitis. Withhold KEYTRUDA for Grade 2; permanently discontinue KEYTRUDA for Grade 3 or 4 or recurrent Grade 2 pneumonitis.
- KEYTRUDA can cause immune-mediated colitis. Colitis occurred in 48 (1.7%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.4%), 3 (1.1%), and 4 (<0.1%) colitis. Monitor patients for signs and symptoms of colitis. Administer corticosteroids for Grade 2 or greater colitis. Withhold KEYTRUDA for Grade 2 or 3; permanently discontinue KEYTRUDA for Grade 4 colitis.
- KEYTRUDA can cause immune-mediated hepatitis. Hepatitis occurred in 19 (0.7%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.1%), 3 (0.4%), and 4 (<0.1%) hepatitis. Monitor patients for changes in liver function. Administer corticosteroids for Grade 2 or greater hepatitis and, based on severity of liver enzyme elevations, withhold or discontinue KEYTRUDA.
- KEYTRUDA can cause hypophysitis. Hypophysitis occurred in 17 (0.6%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.2%), 3 (0.3%), and 4 (<0.1%) hypophysitis. Monitor patients for signs and symptoms of hypophysitis (including hypopituitarism and adrenal insufficiency). Administer corticosteroids and hormone replacement as clinically indicated. Withhold KEYTRUDA for Grade 2; withhold or discontinue for Grade 3 or 4 hypophysitis.
- KEYTRUDA can cause thyroid disorders, including hyperthyroidism, hypothyroidism, and thyroiditis. Hyperthyroidism occurred in 96 (3.4%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.8%) and 3 (0.1%) hyperthyroidism. Hypothyroidism occurred in 237 (8.5%) of 2799 patients receiving KEYTRUDA, including Grade 2 (6.2%) and 3 (0.1%) hypothyroidism. Thyroiditis occurred in 16 (0.6%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.3%) thyroiditis. Monitor patients for changes in thyroid function (at the start of treatment, periodically during treatment, and as indicated based on clinical evaluation) and for clinical signs and symptoms of thyroid disorders. Administer replacement hormones for hypothyroidism and manage hyperthyroidism with thionamides and beta-blockers as appropriate. Withhold or discontinue KEYTRUDA for Grade 3 or 4 hyperthyroidism.
- KEYTRUDA can cause type 1 diabetes mellitus, including diabetic ketoacidosis, which have been reported in 6 (0.2%) of 2799 patients. Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Administer insulin for type 1 diabetes, and withhold KEYTRUDA and administer antihyperglycemics in patients with severe hyperglycemia.

Before prescribing KEYTRUDA, please read the additional Selected Safety Information on the following page and the accompanying [Prescribing Information](#). The [Medication Guide](#) also is available.

KEYTRUDA[®]
(pembrolizumab) Injection 100 mg

SELECTED SAFETY INFORMATION

(continued)

- KEYTRUDA can cause immune-mediated nephritis. Nephritis occurred in 9 (0.3%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.1%), 3 (0.1%), and 4 (<0.1%) nephritis. Monitor patients for changes in renal function. Administer corticosteroids for Grade 2 or greater nephritis. Withhold KEYTRUDA for Grade 2; permanently discontinue KEYTRUDA for Grade 3 or 4 nephritis.
- KEYTRUDA can cause other clinically important immune-mediated adverse reactions. For suspected immune-mediated adverse reactions, ensure adequate evaluation to confirm etiology or exclude other causes. Based on the severity of the adverse reaction, withhold KEYTRUDA and administer corticosteroids. Upon improvement to Grade 1 or less, initiate corticosteroid taper and continue to taper over at least 1 month. Based on limited data from clinical studies in patients whose immune-related adverse reactions could not be controlled with corticosteroid use, administration of other systemic immunosuppressants can be considered. Resume KEYTRUDA when the adverse reaction remains at Grade 1 or less following corticosteroid taper. Permanently discontinue KEYTRUDA for any Grade 3 immune-mediated adverse reaction that recurs and for any life-threatening immune-mediated adverse reaction.
- The following clinically significant immune-mediated adverse reactions occurred in less than 1% (unless otherwise indicated) of 2799 patients: arthritis (1.5%), exfoliative dermatitis, bullous pemphigoid, rash (1.4%), uveitis, myositis, Guillain-Barré syndrome, myasthenia gravis, vasculitis, pancreatitis, hemolytic anemia, and partial seizures arising in a patient with inflammatory foci in brain parenchyma. In addition, myelitis and myocarditis were reported in other clinical trials, including classical Hodgkin lymphoma, and postmarketing use.
- Solid organ transplant rejection has been reported in postmarketing use of KEYTRUDA. Treatment with KEYTRUDA may increase the risk of rejection in solid organ transplant recipients. Consider the benefit of treatment with KEYTRUDA vs the risk of possible organ rejection in these patients.
- KEYTRUDA can cause severe or life-threatening infusion-related reactions, including hypersensitivity and anaphylaxis, which have been reported in 6 (0.2%) of 2799 patients. Monitor patients for signs and symptoms of infusion-related reactions, including rigors, chills, wheezing, pruritus, flushing, rash, hypotension, hypoxemia, and fever. For Grade 3 or 4 reactions, stop infusion and permanently discontinue KEYTRUDA.
- Based on its mechanism of action, KEYTRUDA can cause fetal harm when administered to a pregnant woman. If used during pregnancy, or if the patient becomes pregnant during treatment, apprise the patient of the potential hazard to a fetus. Advise females of reproductive potential to use highly effective contraception during treatment and for 4 months after the last dose of KEYTRUDA.
- In KEYNOTE-006, KEYTRUDA was discontinued due to adverse reactions in 9% of 555 patients with advanced melanoma; adverse reactions leading to discontinuation in more than one patient were colitis (1.4%), autoimmune hepatitis (0.7%), allergic reaction (0.4%), polyneuropathy (0.4%), and cardiac failure (0.4%). Adverse reactions leading to interruption of KEYTRUDA occurred in 21% of patients; the most common ($\geq 1\%$) was diarrhea (2.5%). The most common adverse reactions with KEYTRUDA vs ipilimumab were fatigue (28% vs 28%), diarrhea (26% with KEYTRUDA), rash (24% vs 23%), and nausea (21% with KEYTRUDA). Corresponding incidence rates are listed for ipilimumab only for those adverse reactions that occurred at the same or lower rate than with KEYTRUDA.
- In KEYNOTE-002, KEYTRUDA was discontinued due to adverse reactions in 12% of 357 patients with advanced melanoma; the most common ($\geq 1\%$) were general physical health deterioration (1%), asthenia (1%), dyspnea (1%), pneumonitis (1%), and generalized edema (1%). Adverse reactions leading to interruption of KEYTRUDA occurred in 14% of patients; the most common ($\geq 1\%$) were dyspnea (1%), diarrhea (1%), and maculopapular rash (1%). The most common adverse reactions with KEYTRUDA vs chemotherapy were fatigue (43% with KEYTRUDA), pruritus (28% vs 8%), rash (24% vs 8%), constipation (22% vs 20%), nausea (22% with KEYTRUDA), diarrhea (20% vs 20%), and decreased appetite (20% with KEYTRUDA). Corresponding incidence rates are listed for chemotherapy only for those adverse reactions that occurred at the same or lower rate than with KEYTRUDA.
- It is not known whether KEYTRUDA is excreted in human milk. Because many drugs are excreted in human milk, instruct women to discontinue nursing during treatment with KEYTRUDA and for 4 months after the final dose.

Before prescribing KEYTRUDA, please read the accompanying [Prescribing Information](#). The [Medication Guide](#) also is available.



Merck Oncology

Copyright © 2017 Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc.
All rights reserved. ONCO-1210409-0000 05/17 [keytruda.com](#)

KEYTRUDA[®]
(pembrolizumab) Injection 100 mg